

In the claims:

1-35. (canceled)

36. (Original) A method of stimulating growth of connective tissue or wound healing in a mammal, said method comprising administering to said mammal an effective growth stimulating amount of a polypeptide comprising an amino acid sequence having at least 85% identity with SEQ ID NO:3 or SEQ ID NO:7, or a fragment or analog thereof having the biological activity of PDGF-C, or a polypeptide produced by expression of a polynucleotide comprising a polynucleotide sequence having at least 85% identity with SEQ ID NO:2, SEQ ID NO:4 or SEQ ID NO:6, or of a polynucleotide which hybridizes under stringent conditions with SEQ ID NO:2, SEQ ID NO:4 or SEQ ID NO:6.

37-45. (canceled)

46. (Original) A method of promoting fibroblast mitogenesis in a mammal, comprising the step of administering to said mammal an effective fibroblast mitogenesis promoting amount of a polypeptide comprising an amino acid sequence having at least 85% identity with at least amino acid residues 230 to 345 of SEQ ID NO:3 or of SEQ ID NO:7.

47. (Original) A method of promoting fibroblast mitogenesis in a mammal, comprising administering to said mammal an effective fibroblast mitogenesis promoting amount of a polypeptide comprising an amino acid sequence having at least 85% identity with SEQ ID NO:3 or SEQ ID NO:7, or a fragment or analog thereof having the biological activity of PDGF-C, or a polypeptide produced by expression of a polynucleotide comprising a polynucleotide sequence having at least 85% identity with SEQ ID NO:2, SEQ ID NO:4 or SEQ ID NO:6, or of a polynucleotide which hybridizes under stringent conditions with SEQ ID NO:2, SEQ ID NO:4 or SEQ ID NO:6.

48. (Original) A method of inducing PDGF alpha receptor activation, comprising the step of adding a PDGF alpha-receptor stimulating amount of a polypeptide comprising an amino acid sequence having at least 85% identity with at least amino acid residues 230 to 345 of SEQ ID NO:3 or of SEQ ID NO:7.

49. (Original) A method of inducing PDGF alpha receptor activation, comprising the step of adding a PDGF alpha-receptor stimulating amount of a polypeptide comprising an amino acid sequence having at least 85% identity with SEQ ID NO:3 or SEQ ID NO:7, or a fragment or analog thereof having the biological activity of PDGF-C, or a polypeptide produced by expression of a polynucleotide comprising a polynucleotide sequence having at least 85% identity with SEQ ID NO:2, SEQ ID NO:4 or SEQ ID NO:6, or of a polynucleotide which hybridizes under stringent conditions with SEQ ID NO:2, SEQ ID NO:4 or SEQ ID NO:6.

50-58. (canceled)

59. (Original) A method of promoting angiogenesis in a bird or mammal, said method comprising administering to said bird or mammal an effective angiogenesis promoting amount of a polypeptide comprising a sequence of amino acids having at least 85% identity with at least amino acid residues 230 to 345 of SEQ ID NO:3 or of SEQ ID NO:7.

60. (Original) A method according to claim 59, wherein said polypeptide is administered in the form of a dimer.